



[Science](#)

[Products](#)

[News](#)

[About](#)



[Contact Us](#)



Pfizer Shares Top-Line Results from Phase 2/3 EPIC-PEP Study of PAXLOVID™ for Post-Exposure Prophylactic Use

Friday, April 29, 2022 - 04:30pm



NEW YORK, April 29, 2022 -- [Pfizer Inc.](#) (NYSE: PFE) today shared top-line results from the Phase 2/3 EPIC-PEP (**E**valuation of **P**rotease **I**nhibition for **C**OVID-19 in **P**ost-**E**xposure **P**rophylaxis) study evaluating PAXLOVID™ (nirmatrelvir [PF-07321332] tablets and ritonavir tablets) for post-exposure prophylactic use. In this trial, compared to placebo, Pfizer observed risk reductions of 32% and 37% in adults who received PAXLOVID for five and ten days, respectively, to prevent infection. These results, however, were not statistically significant and, as such, the primary endpoint of reducing the risk of confirmed and symptomatic COVID-19 infection in adults who had been exposed to the virus through a household contact was not met.

safety profile remained generally consistent when PAXLOVID was used for either five or ten days. Analyses of all secondary endpoints and sub-groups are ongoing, and results will be included in the publication or presentation of the final study results.

“We designed the clinical development program for PAXLOVID to be comprehensive and ambitious with the aim of being able to help combat COVID-19 in a very broad population of patients,” said Albert Bourla, Chairman and Chief Executive Officer, Pfizer. “While we are disappointed in the outcome of this particular study, these results do not impact the strong efficacy and safety data we’ve observed in our earlier trial for the treatment of COVID-19 patients at high risk of developing severe illness, and we are pleased to see the growing global use of PAXLOVID in that population.”

PAXLOVID is currently approved or authorized for conditional or emergency use in more than 60 countries across the globe to treat high-risk COVID-19 patients.

Please see Full Emergency Use Authorization (EUA) Prescribing Information available at www.fda.gov and www.COVID19oralRx.com.

About the Phase 2/3 EPIC-PEP Study

The top-line analysis evaluated data from 2,957 adults. Enrolled adults had a negative SARS-CoV-2 rapid antigen test result and were asymptomatic household contacts with exposure within 96 hours to an individual who was symptomatic and recently tested positive for SARS-CoV-2. Each patient was randomized (1:1:1) to receive orally twice daily one of the following: (i) PAXLOVID for five days followed by placebo for 5 days, (ii) PAXLOVID for ten days or (iii) placebo for ten days.

Recruitment began in September 2021 and was completed during the peak of the COVID-19 Omicron wave.

U.S. FDA Emergency Use Authorization Statement

PAXLOVID has not been approved, but has been authorized for emergency use by FDA under an EUA, for the treatment of mild-to-moderate COVID-19 in adults and pediatric patients (12 years of age and older weighing at least 40 kg) with positive results of direct SARS CoV-2 viral testing, and who are at high-risk for progression to severe COVID-19, including hospitalization or death.

The emergency use of PAXLOVID is only authorized for the duration of the declaration that circumstances exist justifying the authorization of the emergency use of drugs and biological products during the COVID-19 pandemic under Section 564(b)(1) of the Act, 21 U.S.C. § 360bbb-3(b)(1), unless the declaration is terminated or authorization revoked sooner.

AUTHORIZED USE

The U.S. Food and Drug Administration (FDA) has issued an Emergency Use Authorization (EUA) for the emergency use of the unapproved product PAXLOVID for the treatment of mild-to-moderate coronavirus disease 2019 (COVID-19) in adults and pediatric patients (12 years of age and older weighing at least 40 kg) with positive results of direct severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) viral testing, and who are at high risk for progression to severe COVID-19, including hospitalization or death.

LIMITATIONS OF AUTHORIZED USE

- PAXLOVID is not authorized for initiation of treatment in patients requiring hospitalization due to severe or critical COVID-19
- PAXLOVID is not authorized for use as pre-exposure or post-exposure prophylaxis for prevention of COVID-19

physician assistants that are licensed or authorized under state law to prescribe drugs in the therapeutic class to which PAXLOVID belongs (i.e., anti-infectives).

PAXLOVID is not approved for any use, including for use for the treatment of COVID-19.

PAXLOVID is authorized only for the duration of the declaration that circumstances exist justifying the authorization of the emergency use of PAXLOVID under 564(b)(1) of the Food Drug and Cosmetic Act unless the authorization is terminated or revoked sooner.

IMPORTANT SAFETY INFORMATION

PAXLOVID is **contraindicated in patients with a history of clinically significant hypersensitivity reactions** (eg, toxic epidermal necrolysis [TEN] or Stevens-Johnson syndrome) to its active ingredients (nirmatrelvir or ritonavir) or any other components of the product.

PAXLOVID is **contraindicated with drugs that are highly dependent on CYP3A** for clearance and for which elevated concentrations are associated with serious and/or life-threatening reactions:

- Alpha1-adrenoreceptor antagonist: alfuzosin
- Analgesics: pethidine, propoxyphene
- Antianginal: ranolazine
- Antiarrhythmic: amiodarone, dronedarone, flecainide, propafenone, quinidine
- Anti-gout: colchicine
- Antipsychotics: lurasidone, pimozide, clozapine

- PDE5 inhibitor: sildenafil (Revatio®) when used for pulmonary arterial hypertension
- Sedative/hypnotics: triazolam, oral midazolam

PAXLOVID is **contraindicated with drugs that are potent CYP3A inducers** where significantly reduced nirmatrelvir or ritonavir plasma concentrations may be associated with the potential for loss of virologic response and possible resistance. PAXLOVID cannot be started immediately after discontinuation of any of the following medications due to the delayed offset of the recently discontinued CYP3A inducer:

- Anticancer drugs: apalutamide
- Anticonvulsant: carbamazepine, phenobarbital, phenytoin
- Antimycobacterials: rifampin
- Herbal Products: St. John's Wort (*hypericum perforatum*)

There are limited clinical data available for PAXLOVID. **Serious and unexpected adverse events may occur** that have not been previously reported with PAXLOVID use.

Risk of Serious Adverse Reactions Due to Drug Interactions: Initiation of PAXLOVID, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving PAXLOVID, may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of PAXLOVID, respectively. These interactions may lead to:

- Clinically significant adverse reactions, potentially leading to severe, life-threatening, or fatal events from greater exposures of concomitant medications

Consult Table 1 of the Fact Sheet for Healthcare Providers for clinically significant drug interactions, including contraindicated drugs. Consider the potential for drug interactions prior to and during PAXLOVID therapy; review concomitant medications during PAXLOVID therapy and monitor for the adverse reactions associated with the concomitant medications.

Hypersensitivity reactions have been reported with PAXLOVID including urticaria, angioedema, dyspnea, mild skin eruptions, and pruritus. Cases of anaphylaxis, TEN, and Stevens-Johnson syndrome have also been reported with components of PAXLOVID (refer to NORVIR labeling). If signs and symptoms of a clinically significant hypersensitivity reaction or anaphylaxis occur, immediately discontinue PAXLOVID and initiate appropriate medications and/or supportive care.

Hepatotoxicity: Hepatic transaminase elevations, clinical hepatitis, and jaundice have occurred in patients receiving ritonavir. Therefore, caution should be exercised when administering PAXLOVID to patients with **pre-existing liver diseases, liver enzyme abnormalities, or hepatitis.**

Because nirmatrelvir is co-administered with ritonavir, there may be a **risk of HIV-1 developing resistance to HIV protease inhibitors** in individuals with uncontrolled or undiagnosed HIV-1 infection.

Adverse events in the PAXLOVID group ($\geq 1\%$) that occurred at a greater frequency (≥ 5 subject difference) than in the placebo group were dysgeusia (6% and $< 1\%$, respectively), diarrhea (3% and 2%), hypertension (1% and $< 1\%$), and myalgia (1% and $< 1\%$). The proportions of subjects who discontinued treatment due to an adverse event were 2% in the PAXLOVID group and 4% in the placebo group.

The following adverse reactions have been identified during post-authorization use of PAXLOVID. Because these

Immune System Disorders: Hypersensitivity reactions

Required Reporting for Serious Adverse Events and Medication Errors: The prescribing healthcare provider and/or the provider's designee is/are responsible for mandatory reporting of all serious adverse events and medication errors potentially related to PAXLOVID within 7 calendar days from the healthcare provider's awareness of the event.

Submit adverse event and medication error reports to FDA MedWatch using one of the following methods:

- **Online:** <https://www.fda.gov/medwatch/report.htm>
- **Complete and submit a postage-paid FDA Form 3500 and returning by mail/fax**
- **Call 1-800-FDA-1088 to request a reporting form**

In addition, please provide a copy of all FDA MedWatch forms to: <http://www.pfizersafetyreporting.com/> or by fax (1-866-635-8337) or phone (1-800-438-1985).

PAXLOVID is an inhibitor of CYP3A and may increase plasma concentrations of drugs that are primarily metabolized by CYP3A. Co-administration of PAXLOVID with drugs highly dependent on CYP3A for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events is contraindicated. Co-administration with other CYP3A substrates may require a dose adjustment or additional monitoring.

Nirmatrelvir and ritonavir are CYP3A substrates; therefore, drugs that induce CYP3A may decrease nirmatrelvir and ritonavir plasma concentrations and reduce PAXLOVID therapeutic effect.

Pregnancy: There are no available human data on the use of nirmatrelvir during pregnancy to evaluate for a

studies with ritonavir are insufficient to identify a drug-associated risk of miscarriage. There are maternal and fetal risks associated with untreated COVID-19 in pregnancy.

Lactation: There are no available data on the presence of nirmatrelvir in human or animal milk, the effects on the breastfed infant, or the effects on milk production. A transient decrease in body weight was observed in the nursing offspring of rats administered nirmatrelvir. Limited published data reports that ritonavir is present in human milk. There is no information on the effects of ritonavir on the breastfed infant or the effects of the drug on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for PAXLOVID and any potential adverse effects on the breastfed infant from PAXLOVID or from the underlying maternal condition. Breastfeeding individuals with COVID-19 should follow practices according to clinical guidelines to avoid exposing the infant to COVID-19.

Contraception: Use of ritonavir may reduce the efficacy of combined hormonal contraceptives. Advise patients using combined hormonal contraceptives to use an effective alternative contraceptive method or an additional barrier method of contraception.

Pediatrics: PAXLOVID is not authorized for use in pediatric patients younger than 12 years of age or weighing less than 40 kg. The safety and effectiveness of PAXLOVID have not been established in pediatric patients. The authorized adult dosing regimen is expected to result in comparable serum exposures of nirmatrelvir and ritonavir in patients 12 years of age and older and weighing at least 40 kg as observed in adults, and adults with similar body weight were included in the trial EPIC-HR.

Systemic exposure of nirmatrelvir increases in renally impaired patients with increase in the severity of renal impairment.

patients about renal dosing instructions. **PAXLOVID is not recommended in patients with severe renal impairment** (eGFR <30 mL/min based on CKD-EPI formula) until more data are available; the appropriate dosage for patients with severe renal impairment has not been determined.

No dosage adjustment of PAXLOVID is needed for patients with either mild (Child-Pugh Class A) or moderate (Child-Pugh Class B) hepatic impairment. No pharmacokinetic or safety data are available regarding the use of nirmatrelvir or ritonavir in subjects with severe hepatic impairment (Child-Pugh Class C); therefore, **PAXLOVID is not recommended for use in patients with severe hepatic impairment.**

About Pfizer: Breakthroughs That Change Patients' Lives

At Pfizer, we apply science and our global resources to bring therapies to people that extend and significantly improve their lives. We strive to set the standard for quality, safety and value in the discovery, development and manufacture of health care products, including innovative medicines and vaccines. Every day, Pfizer colleagues work across developed and emerging markets to advance wellness, prevention, treatments and cures that challenge the most feared diseases of our time. Consistent with our responsibility as one of the world's premier innovative biopharmaceutical companies, we collaborate with health care providers, governments and local communities to support and expand access to reliable, affordable health care around the world. For more than 170 years, we have worked to make a difference for all who rely on us. We routinely post information that may be important to investors on our website at www.Pfizer.com. In addition, to learn more, please visit us on www.Pfizer.com and follow us on Twitter at [@Pfizer](#) and [@Pfizer News](#), [LinkedIn](#), [YouTube](#) and like us on Facebook at Facebook.com/Pfizer.

Disclosure Notice

[Science](#)[Products](#)[News](#)[About](#)[Contact Us](#)

This release contains forward-looking information about Pfizer's efforts to combat COVID-19 and PAXLOVID (including qualitative assessments of available data, potential benefits, expectations for clinical trials, the anticipated timing of data readouts, regulatory submissions, regulatory approvals or authorizations and use in high risk COVID-19 patients), involving substantial risks and uncertainties that could cause actual results to differ materially from those expressed or implied by such statements. Risks and uncertainties include, among other things, the uncertainties inherent in research and development, including the ability to meet anticipated clinical endpoints, commencement and/or completion dates for clinical trials, regulatory submission dates, regulatory approval dates and/or launch dates, as well as risks associated with preclinical and clinical data, including the possibility of unfavorable new preclinical, clinical or safety data and further analyses of existing preclinical, clinical or safety data, including the risk that final results from EPIC-SR could differ from the interim data; the ability to produce comparable clinical or other results including efficacy, safety and tolerability profile observed to date, in additional studies or in larger, more diverse populations following commercialization; uncertainties regarding the commercial impact of the results of the EPIC-PEP trial; the ability of PAXLOVID to maintain efficacy against emerging virus variants; the risk that serious and unexpected adverse events may occur that have not been previously reported with PAXLOVID use; the risk that preclinical and clinical trial data are subject to differing interpretations and assessments, including during the peer review/publication process, in the scientific community generally, and by regulatory authorities; whether regulatory authorities will be satisfied with the design of and results from these and any future preclinical and clinical studies; whether and when any drug applications or submissions to request emergency use or conditional marketing authorization for any potential indications for PAXLOVID may be filed in particular jurisdictions and if obtained, whether or when such emergency use authorization or licenses will expire or terminate; whether and when regulatory authorities in any jurisdictions may approve any applications or submissions for PAXLOVID that may be pending or filed (including a potential new drug application submission in the U.S. and submissions in other jurisdictions), which will depend on myriad factors, including making a determination as to whether



processes, safety and/or other matters that could affect the availability or commercial potential of PAXLOVID, including development of products or therapies by other companies; risks related to the availability of raw materials for PAXLOVID; the risk that we may not be able to create or scale up manufacturing capacity on a timely basis or maintain access to logistics or supply channels commensurate with global demand, which would negatively impact our ability to supply the estimated numbers of courses of PAXLOVID within the projected time periods; whether and when additional purchase agreements will be reached; the risk that demand for any products may be reduced or no longer exist; uncertainties regarding the impact of COVID-19 on Pfizer's business, operations and financial results; and competitive developments.

A further description of risks and uncertainties can be found in Pfizer's Annual Report on Form 10-K for the fiscal year ended December 31, 2021 and in its subsequent reports on Form 10-Q, including in the sections thereof captioned "Risk Factors" and "Forward-Looking Information and Factors That May Affect Future Results", as well as in its subsequent reports on Form 8-K, all of which are filed with the U.S. Securities and Exchange Commission and available at www.sec.gov and www.pfizer.com.

Medicines

Pfizer Contacts:

Media Relations

+1 (212) 733-1226

IR@pfizer.com

Related press releases

04.27.2022

[Pfizer and Biohaven's VYDURA® \(Rimegepant\) Granted First Ever Marketing Authorization by European Commission for Both Acute Treatment of Migraine and Prophylaxis of Episodic Migraine](#)

Medicines

04.22.2022

[Pfizer Voluntary Nationwide Recall of Lots of ACCUPRIL® \(Quinapril HCl\) Due to N-Nitroso-Quinapril Content](#)

Medicines



[Science](#)

[Products](#)

[News](#)

[About](#)

[Contact Us](#)

04.12.2022

Myovant Sciences and Pfizer Provide Update on Supplemental New Drug Application for MYFEMBREE® for the Management of Moderate to Severe Pain Associated with Endometriosis

Medicines

[See More Press Releases](#)



[Investors](#)

[Careers](#)

[Grant Seekers](#)

[Healthcare Professionals](#)

[Privacy Statement](#)

[Terms of Use](#)



[Science](#)

[Products](#)

[News](#)

[About](#)

[Contact Us](#)

United States © 2022 Pfizer Inc. All rights reserved

